

S51.05

Treatment-seeking gamblers and Parkinson's disease: Case reports

O. Simon. *Center for Pathological Gambling, Community Psychiatry Services, Department of Adult Psychiatry, University Hospital of Lausanne, Lausanne, Switzerland*

Pathological gambling (PG) is a rare but well-established behavioural disorders of Parkinson's disease (PD) patients treated with dopamine agonist. We investigate the chronological relationship between PD and diagnosis of PG within treatment-seeking gamblers.

Sample and Methods: From 174 consecutively admitted pathological gamblers we identified 4 patients with PD. Standardized medical records include socio-demographic characteristics, past gaming behaviours and gambling-treatment modalities as well as the evolution of PD and the pro-dopaminergic medication history.

Results: All four patients developed PG after the onset of PD treatment. Three patients described a period of chronic exposure to gaming facilities years before and after onset of pro-dopaminergic treatment, with occasional or regular gambling, but without a compulsive component. In one patient, PG appeared suddenly without any previous gaming behaviour after the onset of medication. Despite a high treatment motivation, gambling specific cognitive therapy was unsuccessful. (c) Implication for the Field

The relationship between PD and PG appears to be complex: Confounders such as psychosocial factors or increase in accessibility of gambling opportunities may account for these findings. This case series confirm that pro-dopaminergic treatment can induce PG behaviour, but may be more likely in "at risk" groups. These patients pose specific treatment challenges.

Symposium: The role of 5-HT1AR in pathophysiology and treatment of schizophrenia

S48.01

The postsynaptic 5-HT1A-receptor and its role for cognitive functions in mice

B. Bert, J. Rothe, H. Fink. *Freie Universität Berlin, School for Veterinary Medicine, Institute of Pharmacology and Toxicology, Berlin, Germany*

Schizophrenia is often accompanied by cognitive dysfunctions. Post-mortem and in vivo studies have revealed increased cortical 5-HT1A-receptor density and it was assumed that 5-HT1A-receptor active drugs could enhance cognition. Moreover, partial 5-HT1A-receptor agonists positively affected verbal memory in schizophrenic patients. However, the role of the 5-HT1A-receptor for cognition has not been fully clarified.

Recently, we have introduced transgenic mice overexpressing the 5-HT1A-receptor postsynaptically in the cortex and hippocampus. Function of the surplus receptors was verified by receptor activation with the agonist 8-OH-DPAT.

In this study we further investigated the role of postsynaptic 5-HT1A-receptors for cognition. Therefore, our mice were tested in the inhibitory avoidance, Morris water maze, and hole-board habituation task. Moreover, the effects of low and high doses of 8-OH-DPAT were examined in the inhibitory avoidance task.

Our transgenic mice showed no overall cognitive deficit. As a tendency, inhibitory avoidance retention was impaired in transgenic mice compared to wild-type controls. Both genotypes showed similar spatial learning abilities in the Morris water maze and habituated to the hole-board in a comparable manner. Anterograde amnesia induced by 8-OH-DPAT was in transgenic mice already apparent in a third of the dose used for wild-type mice. Retrograde amnesia could not be triggered.

Since the transgenic mice show untreated a rather normal behaviour, we assume that they possess compensatory mechanisms. However, after activation of the postsynaptic 5-HT1A-receptors the differences between wild-type and transgenic mice became more clear. Hence, our findings suggest that the cortical and hippocampal 5-HT1A-receptors play rather a modulatory role in learning.

S48.02

Effect of Serotonin-1A receptor on behavioral changes in animal model of schizophrenia-like behavior

V. Bubenikova-Valesova¹, M. Votava², T. Palenicek¹, J. Horacek^{1,2}, C. Hoschl^{1,2}. ¹Prague Psychiatric Center, Prague, Czech Republic ²3rd Faculty of Medicine Charles University, Prague, Czech Republic

Some antipsychotics act as partial agonists on serotonin-1A receptors (5-HT1AR) localized postsynaptically in cortex and hippocampus and presynaptically on the serotonergic cell bodies and dendrites in raphe nuclei.

Our study's aim was to investigate the effect of pre- and postsynaptic 5-HT1AR activation on MK-801 (0.1, 0.3 mg/kg)-induced sensorimotor gating deficits and hyperlocomotion in a rat model of schizophrenia-like behavior. To investigate the effect of presynaptic receptor activation we used a partial agonist (buspirone; 1,10 mg/kg) and a low dose of full agonist (8-OH-DPAT; 0.025 mg/kg). The effect on both pre- and postsynaptic receptors was investigated by a high dose of full agonist (8-OH-DPAT; 1 mg/kg).

We found that buspirone in both doses had no effect on MK-801-induced deficit in sensorimotor gating. Contrarily, the low dose of 8-OH-DPAT ameliorated the deficit. The MK-801-induced hyperlocomotion was decreased by buspirone as well as by the low dose of 8-OH-DPAT. Activation of both pre- and postsynaptic 5-HT1AR had an opposite effect on MK-801-induced behavior.

Our findings accord with the published results that partial 5-HT1AR agonists could be effective in schizophrenia treatment, but full potent agonists could exacerbate psychotic symptoms. Observed differences between buspirone and the low dose of 8-OH-DPAT could be due to inhibition of D2 receptor.

This research was supported by grant MZ0PCP2005 from the MZCR and by the projects 1M0517 from the MSMT

S48.03

The influence of polymorphism for Serotonin 5HT-1A receptor on phenotypic variables in schizophrenia

J. Horacek^{1,2,3}, F. Spaniel², M. Brunovsky³, T. Novak², B. Tislerova², M. Kopecek³, P. Mohr¹, C. Hoschl^{1,3}. ¹Prague Psychiatric Center, Prague, Czech Republic ²3rd Faculty of Medicine, Charles University, Prague, Czech Republic ³Centre of Neuropsychiatric Studies, Prague, Czech Republic

Dysfunction of the serotonin system has been implicated in schizophrenia. 5-HT1A and 5-HT2A serotonin receptors are involved in